

Culture of microalgae with ultrafiltered seawater: from a feasibility study to an industrial development

Clémence Cordier ^{a*}, Christophe Stavrakakis ^{b*}, Adeline Lange ^c, Frédéric Chenier ^c, Philippe Moulin ^{a*}

^a Laboratoire de Mécanique, Modélisation et Procédés Propres (M2P2-CNRS-UMR 7340), Aix-Marseille Université, Europôle de l'Arbois, BP 80, Bat. Laennec, Hall C, 13545 Aix-en-Provence cedex 04, France

^b Plateforme expérimentale Mollusques Marins, Station Ifremer de Bouin, Polder des Champs, 85230 Bouin, France

^c Vendée Naissain, Polder des Champs, 85230 Bouin, France

CONTEXT

The culture of microalgae is essential for the production and maintenance of bivalves. One of the major challenges is to reduce cultivation costs while maintaining the reliability of microalgae forages over long time. The aim of this work is to use ultrafiltered (UF) seawater to cultivate them.

MATERIAL AND METHODS

Membranes: Aquasource hollow fibre PES membranes (0.02 µm), in-out configuration.

Pilot: Semi industrial unit, completely automated, able to treat 20 m³.d⁻¹.

Microalgae cultivation: Every day a volume of microalgae was withdrawn to feed oysters + refill of tanks with water


• Daily monitoring of: **microalgae concentration** + **water quality** (temperature, pH, salinity, O₂) and **microscopic observations**



T-ISOCHRYSIS

T-isochrysis concentrations are higher in UF seawater than control seawater: the gain is from 6 % to 30 % over the 5 experiments.


→ Ultrafiltered water is adapted to cultivation of *T-isochrysis*




Evolution of *T-isochrysis* concentration vs. time

COMPARISON WITH BOREHOLE WATER

UF SEAWATER BOREHOLE WATER




Comparison of *T-iso* cultivation in a shellfish hatchery in batch culture with UF seawater (UFSW) and borehole water (BW)




UF: greater mobility of the cultures and less contaminations



→ Confirmation that UF seawater is suitable for microalgae production



Appearance, mobility, and contamination of the 30 L UF and BW. Bleu = good, orange = average and yellow = crash of culture



Average growth curve of *T-Iso* for the two replicates and the two types of water in a 2 L flask




T = 20 °C, V = 300 L, bubling air + CO₂, light

Microalgae cultivation - Water treatment steps

TETRASELMIS

Tetraselmis concentrations are similar in both water qualities

→ Ultrafiltered water is adapted to cultivation of *Tetraselmis*



Evolution of *Tetraselmis* concentration vs. time

Parasites were observed in microalgae cultivated in control seawater but none in UF seawater

→ A protection of *Tetraselmis* toward parasites is obtained with ultrafiltration



Microscopic observations in control seawater

CONCLUSION

Ultrafiltered water has shown **efficiency for microalgae cultivation** with **rapid growth** and/or significant **reduction in contamination** compared to cultivation with control waters. These conclusions and the **ease of recovering water** (linked to the reduction in treatment stages) allowed a **transfer of technology**: the 300 L cultures carried out on the experimental platform (Ifremer) are now produced with ultrafiltered water since early 2019..